arXiv:1601.05515v1 [math.NT] 21 Jan 2016

On the Seventh Power Moment of $\Delta(x)$

Jinjiang Li

Department of Mathematics,
China University of Mining and Technology,
Beijing 100083, P. R. China
Email: jinjiang.li.math@gmail.com

Abstract: Let $\Delta(x)$ be the error term of the Dirichlet divisor problem. In this paper, we establish an asymptotic formula of the seventh-power moment of $\Delta(x)$ and prove that

$$\int_2^T \Delta^7(x)\mathrm{d}x = \frac{7(5s_{7;3}(d) - 3s_{7;2}(d) - s_{7;1}(d))}{2816\pi^7}T^{11/4} + O(T^{11/4-\delta_7+\varepsilon})$$

with $\delta_7 = 1/336$, which improves the previous result.

Keywords: Dirichlet divisor problem; higher-power moment; asymptotic formula.

1 Introduction and main result

Throughout this paper, let $d(n)$ denote the Dirichlet divisor function. In 1838, Dirichlet proved that the error term

$$\Delta(x) := \sum_{n\leqslant x} d(n) - x\log x - (2\gamma - 1)x, \qquad \text{with } x \geqslant 2,$$

satisfies $\Delta(x) \ll x^{1/2}$. Here γ is Euler's constant. Since then the determination of the exact order of $\Delta(x)$ has been called Dirichlet's divisor problem. Many writers have sharpened Dirichlet's bound for $\Delta(x)$. The latest result is due to Huxley [5], who proved that

$$\Delta(x) \ll x^{131/416}(\log x)^{26957/8320}.$$

For a survey of the history of this problem, see Krätzel [10].

In the opposite direction, Hardy [3] proved that

$$\Delta(x) = \begin{cases} \Omega_+\left(x^{1/4}(\log x)^{1/4}\log\log x\right), \\ \Omega_-\left(x^{1/4}\right). \end{cases}$$

The best results in this direction to date are

$$\Delta(x) = \Omega_+\left(x^{1/4}(\log x)^{1/4}(\log\log x)^{(3+\log 4)/4}\exp(-c\sqrt{\log\log\log x})\right)$$

and

$$\Delta(x) = \Omega_- \left(x^{1/4} \exp(c(\log\log x)^{1/4}(\log\log\log x)^{-3/4} \right)$$

for some constant $c > 0$, due to Hafner [2] and [1] respectively. It is conjectured that $\Delta(x) \ll x^{1/4+\varepsilon}$ is true for every $\varepsilon > 0$. The evidence in support of this conjecture has been given by Tong [14] and Ivic [6], who proved, respectively, that

$$\int_2^T \Delta^2(x)\mathrm{d}x = \frac{\zeta^4(\frac{3}{2})}{6\pi^2\zeta(3)} T^{3/2} + O(T\log^5 T) \tag{1.1}$$

and

$$\int_2^T |\Delta(x)|^A \mathrm{d}x \ll_\varepsilon T^{1+\frac{A}{4}+\varepsilon}, \qquad \text{for } 0 \leqslant A \leqslant \frac{35}{4} \tag{1.2}$$

and any $\varepsilon > 0$. On the other hand , Voronoï [16] proved that

$$\int_2^T \Delta(x)\mathrm{d}x = T/4 + O(T^{3/4}), \tag{1.3}$$

which in conjunction with (1.1) and (1.2) shows that $\Delta(x)$ has a lot of sign changes and cancellations between the positive and negative portions.

Tsang [15] first studied the third and fourth-power moments of $\Delta(x)$. He proved that

$$\int_2^T \Delta^3(x)\mathrm{d}x = \frac{3c_1}{28\pi^3} T^{7/4} + O(T^{7/4-\delta_3+\varepsilon}), \tag{1.4}$$

$$\int_2^T \Delta^4(x)\mathrm{d}x = \frac{3c_2}{64\pi^4} T^2 + O(T^{2-\delta_4+\varepsilon}), \tag{1.5}$$

where $\delta_3 = 1/14, \delta_4 = 1/23$ and

$$c_1 := \sum_{\alpha,\beta,h\in\mathbb{N}} (\alpha\beta(\alpha+\beta))^{-3/2} h^{-9/4} |\mu(h)| d(\alpha^2 h) d(\beta^2 h) d((\alpha+\beta)^2 h),$$

$$c_2 := \sum_{\substack{n,m,k,\ell\in\mathbb{N} \\ \sqrt{n}+\sqrt{m}=\sqrt{k}+\sqrt{\ell}}} (nmk\ell)^{-3/4} d(n)d(m)d(k)d(\ell),$$

and $\mu(h)$ is the Möbius function.

In [18], Zhai proved that (1.4) holds for $\delta_3 = 1/4$. Ivić and Sargos [8] proved that (1.4) holds for $\delta_3 = 7/20$. Following the approach of Tsang [15], Zhai [18] proved that the equation (1.5) holds for $\delta_4 = 2/41$. This approach used the method of exponential sums. In particular, if the exponent pair conjecture is true, namely, if $(\varepsilon, 1/2+\varepsilon)$ is an exponent pair, then the equation (1.5) holds for $\delta_4 = 1/14$. Moreover, in [8], Ivić and Sargos proved a substantially better result that the equation (1.5) holds for $\delta_4 = 1/12$. Later, combining the method of [8] and a deep result of Robert and Sargos [12], Zhai [20] proved that the equation (1.5) holds for $\delta_4 = 3/28$. Recently, Kong [9] proved that $\delta_4 = 1/8$.

By a unified approach, Zhai [19] proved that the asymptotic formula

$$\int_1^T \Delta^k(x)\mathrm{d}x = C_k T^{1+k/4} + O\left(T^{1+k/4-\delta_k+\varepsilon}\right) \tag{1.6}$$

holds for $3 \leqslant k \leqslant 9$, where C_k and $0 < \delta_k < 1$ are explicit constants. He gives $\delta_5 = 1/64, \delta_6 = 35/4742, \delta_7 = 17/6312, \delta_8 = 8/9433, \delta_9 = 13/75216$. The asymptotic formula (1.6) improved the result of Heath-Brown [4]. When $k = 5$, the asymptotic formula (1.6) holds for $\delta_5 = 1/64$, which improved an earlier exponent $\delta_5 = 5/816$ proved in [18] by the approach of Tsang [15]. In [21], Zhang and Zhai improved the previous result of the case $k = 5$ and proved $\delta_5 = 3/80$. Meanwhile, Wang [17] studied the case $k = 6$ and proved $\delta_6 = 3/248$, which improved the result of Zhai [19], i.e. $\delta_6 = 35/4742$.

The aim of this paper is to improve the value of $\delta_7 = 17/6312$, which is achieved by Zhai [19]. The main result is the following

Theorem 1.1 *We have*

$$\int_2^T \Delta^7(x)\mathrm{d}x = \frac{7(5s_{7;3}(d) - 3s_{7;2}(d) - s_{7;1}(d))}{2816\pi^7} T^{11/4} + O(T^{11/4-\delta_7+\varepsilon})$$

with $\delta_7 = 1/336$, *where*

$$s_{7;3}(d) = \sum_{\substack{n,m,k,\ell,r,s,q\in\mathbb{N}^* \\ \sqrt{n}+\sqrt{m}+\sqrt{k}+\sqrt{\ell}=\sqrt{r}+\sqrt{s}+\sqrt{q}}} \frac{d(n)d(m)d(k)d(\ell)d(r)d(s)d(q)}{(nmk\ell rsq)^{3/4}},$$

$$s_{7;2}(d) = \sum_{\substack{n,m,k,\ell,r,s,q\in\mathbb{N}^* \\ \sqrt{n}+\sqrt{m}+\sqrt{k}+\sqrt{\ell}+\sqrt{r}=\sqrt{s}+\sqrt{q}}} \frac{d(n)d(m)d(k)d(\ell)d(r)d(s)d(q)}{(nmk\ell rsq)^{3/4}},$$

$$s_{7;1}(d) = \sum_{\substack{n,m,k,\ell,r,s,q\in\mathbb{N}^* \\ \sqrt{n}+\sqrt{m}+\sqrt{k}+\sqrt{\ell}+\sqrt{r}+\sqrt{s}=\sqrt{q}}} \frac{d(n)d(m)d(k)d(\ell)d(r)d(s)d(q)}{(nmk\ell rsq)^{3/4}}.$$

Notations. Throughout this paper, $\|x\|$ denotes the distance from x to the nearest integer, i.e., $\|x\| = \min_{n\in\mathbb{Z}} |x - n|$. $[x]$ denotes the integer part of x; $n \sim N$ means $N < n \leqslant 2N$; $n \asymp N$ means $C_1 N \leqslant n \leqslant C_2 N$ with positive constants C_1, C_2 satisfying $C_1 < C_2$. ε always denotes an arbitrary small positive constant which may not be the same at different occurances. We shall use the estimates $d(n) \ll n^\varepsilon$. Suppose $f : \mathbb{N} \to \mathbb{R}$ is any function satisfying $f(n) \ll n^\varepsilon$, $k \geqslant 2$ is a fixed integer. Define

$$s_{k;\ell}(f) := \sum_{\substack{n_1,\cdots,n_\ell,n_{\ell+1},\cdots,n_k\in\mathbb{N}^* \\ \sqrt{n_1}+\cdots+\sqrt{n_\ell}=\sqrt{n_{\ell+1}}+\cdots+\sqrt{n_k}}} \frac{f(n_1)f(n_2)\cdots f(n_k)}{(n_1n_2\cdots n_k)^{3/4}}, \quad 1 \leqslant \ell < k. \tag{1.7}$$

We shall use $s_{k;\ell}(f)$ to denote both of the series (1.7) and its value. Suppose $y > 1$ is a large parameter, and we define

$$s_{k;\ell}(f;y) := \sum_{\substack{n_1,\cdots,n_\ell,n_{\ell+1},\cdots,n_k\leqslant y \\ \sqrt{n_1}+\cdots+\sqrt{n_\ell}=\sqrt{n_{\ell+1}}+\cdots+\sqrt{n_k}}} \frac{f(n_1)f(n_2)\cdots f(n_k)}{(n_1n_2\cdots n_k)^{3/4}}, \quad 1 \leqslant \ell < k.$$

2 Preliminary Lemmas

Lemma 2.1 *Suppose* $k \geqslant 3, (i_1, i_2, \cdots, i_{k-1}) \in \{0,1\}^{k-1}$ *such that*

$$\sqrt{n_1} + (-1)^{i_1}\sqrt{n_2} + (-1)^{i_2}\sqrt{n_3} + \cdots + (-1)^{i_{k-1}}\sqrt{n_k} \neq 0.$$

Then we have

$$|\sqrt{n_1}+(-1)^{i_1}\sqrt{n_2}+(-1)^{i_2}\sqrt{n_3}+\cdots+(-1)^{i_{k-1}}\sqrt{n_k}|\gg \max(n_1,n_2,\cdots,n_k)^{-(2^{k-2}-2^{-1})}.$$

Proof. See Lemma 2.2 of [19].

Lemma 2.2 *If $g(x)$ and $h(x)$ are continuous real-valued functions of x and $g(x)$ is monotonic, then*

$$\int_a^b g(x)h(x)\mathrm{d}x \ll \left(\max_{a\leqslant x\leqslant b}|g(x)|\right)\left(\max_{a\leqslant u<v\leqslant b}\left|\int_u^v h(x)\mathrm{d}x\right|\right).$$

Proof. See Lemma 1 of [15].

Lemma 2.3 *Suppose $A,B\in\mathbb{R}, A\neq 0$. Then*

$$\int_T^{2T}\cos(A\sqrt{t}+B)\mathrm{d}t\ll T^{1/2}|A|^{-1}.$$

Proof. It follows from Lemma 2.2 easily.

Lemma 2.4 *Suppose $K\geqslant 10, \alpha,\beta\in\mathbb{R}, 2K^{-1/2}\leqslant|\alpha|\ll K^{1/2}$ and $0<\delta<1/2$. Then we have*

$$\#\{k\sim K:\|\beta+\alpha\sqrt{t}\|<\delta\}\ll K\delta+K^{1/2+\varepsilon}.$$

Proof. See Lemma 4 of [20].

Lemma 2.5 *Let a,δ be real numbers, $0<\delta<a/4$, and let k be a positive integer. There exists a function $\varphi(y)$ which is k times continuously differentiable and such that*

$$\begin{cases} \varphi(y)=1, & \text{for} \quad |y|\leqslant a-\delta,\\ 0<\varphi(y)<1, & \text{for} \quad a-\delta<|y|<a+\delta,\\ \varphi(y)=0, & \text{for} \quad |y|\geqslant a+\delta, \end{cases}$$

and its Fourier transform

$$\Phi(x)=\int_{-\infty}^{+\infty}e(-xy)\varphi(y)\mathrm{d}y$$

satisfies the inequality

$$|\Phi(x)|\leqslant\min\left(2a,\frac{1}{\pi|x|},\frac{1}{\pi|x|}\left(\frac{k}{2\pi|x|\delta}\right)^k\right).$$

Proof. See [11] or [13].

Lemma 2.6 *Let $d(n)$ denote the divisor function. Then we have*

$$|s_{k;\ell}(d)-s_{k;\ell}(d;y)|\ll y^{-1/2+\varepsilon},\qquad 1\leqslant\ell<k.$$

Proof. See Lemma 3.1 of [19].

Lemma 2.7 *Suppose* $k \geqslant 3; (i_1, i_2, \cdots, i_{k-1}) \in \{0,1\}^{k-1}, (i_1, i_2, \cdots, i_{k-1}) \neq (0, 0, \cdots, 0)$; $1 < N_1, N_2, \cdots, N_k; 0 < \Delta \ll E^{1/2}, E = \max(N_1, N_2, \cdots, N_k)$. *Let*

$$\mathscr{A} = \mathscr{A}(N_1, N_2, \cdots, N_k; i_1, i_2, \cdots, i_k; \Delta)$$

denote the number of solutions of the inequality

$$|\sqrt{n_1} + (-1)^{i_1}\sqrt{n_2} + (-1)^{i_2}\sqrt{n_3} + \cdots + (-1)^{i_{k-1}}\sqrt{n_k}| < \Delta$$

with $n_j \sim N_j, j = 1, 2, \cdots, k$. *Then*

$$\mathscr{A} \ll \Delta E^{-1/2} N_1 N_2 \cdots N_k + E^{-1} N_1 N_2 \cdots N_k.$$

Proof. See Lemma 2.4 of [19].

Lemma 2.8 *Suppose* $1 \leqslant N \leqslant M \leqslant K \leqslant L, 1 \leqslant R \leqslant S \leqslant Q, L \asymp Q$ *and* $0 < \Delta \ll Q^{1/2}$. *Let* $\mathscr{A}_1(N, M, K, L, R, S, Q; \Delta)$ *denote the number of solutions of the inequality*

$$0 < |\sqrt{n} + \sqrt{m} + \sqrt{k} + \sqrt{\ell} - \sqrt{r} - \sqrt{s} - \sqrt{q}| < \Delta \tag{2.1}$$

with $n \sim N, m \sim M, k \sim K, \ell \sim L, r \sim R, s \sim S, q \sim Q$. *Then*

$$\mathscr{A}_1(N, M, K, L, R, S, Q; \Delta) \ll \Delta Q^{1/2} NMKLRS + NMKRSL^{1/2+\varepsilon}.$$

In particular, if $\Delta Q^{1/2} \gg 1$, *then*

$$\mathscr{A}_1(N, M, K, L, R, S, Q; \Delta) \ll \Delta Q^{1/2} NMKLRS.$$

Proof. If $(n, m, k, \ell.r, s, q)$ satisfies (2.1), then

$$(\sqrt{n} + \sqrt{m} + \sqrt{k} - \sqrt{r} - \sqrt{s}) + \sqrt{\ell} = \sqrt{q} + \theta\Delta$$

for some $0 < |\theta| < 1$. Thus, we have

$$2\ell^{1/2}(\sqrt{n} + \sqrt{m} + \sqrt{k} - \sqrt{r} - \sqrt{s}) + (\sqrt{n} + \sqrt{m} + \sqrt{k} - \sqrt{r} - \sqrt{s})^2 + \ell = q + u$$

with $|u| = |2q^{1/2}\theta\Delta + \theta^2\Delta^2| \leqslant 2q^{1/2}\Delta + \Delta^2 \ll \Delta Q^{1/2}$. Then we have

$$q = 2\ell^{1/2}(\sqrt{n} + \sqrt{m} + \sqrt{k} - \sqrt{r} - \sqrt{s}) + (\sqrt{n} + \sqrt{m} + \sqrt{k} - \sqrt{r} - \sqrt{s})^2 + \ell - u$$

with $|u| \leqslant C\Delta Q^{1/2}$ for some absolute positive constant $C > 0$. Hence the quantity of $\mathscr{A}_1(N, M, K, L, R, S, Q; \Delta)$ does not exceed the number of solutions of

$$\left|2\ell^{1/2}(\sqrt{n} + \sqrt{m} + \sqrt{k} - \sqrt{r} - \sqrt{s}) + (\sqrt{n} + \sqrt{m} + \sqrt{k} - \sqrt{r} - \sqrt{s})^2 + \ell - q\right| < C\Delta Q^{1/2} \tag{2.2}$$

with $n \sim N, m \sim M, k \sim K, \ell \sim L, r \sim R, s \sim S, q \sim Q$.

If $\Delta Q^{1/2} \gg 1$, then for fixed (n, m, k, ℓ, r, s), the number of q for which (2.2) holds is $\ll 1 + \Delta Q^{1/2} \ll \Delta Q^{1/2}$. Hence

$$\mathscr{A}_1(N, M, K, L, R, S, Q; \Delta) \ll \Delta Q^{1/2} NMKLRS.$$

Now suppose $\Delta Q^{1/2} \leqslant 1/4C$. Then for fixed (n, m, k, ℓ, r, s), there is at most one q such that (2.2) holds. If such q exists, then we have

$$\left\|2\ell^{1/2}(\sqrt{n}+\sqrt{m}+\sqrt{k}-\sqrt{r}-\sqrt{s})+(\sqrt{n}+\sqrt{m}+\sqrt{k}-\sqrt{r}-\sqrt{s})^2\right\| < C\Delta Q^{1/2}. \tag{2.3}$$

We shall use Lemma 2.4 to bound the number of solutions of (2.3) with $\alpha = 2(\sqrt{n}+\sqrt{m}+\sqrt{k}-\sqrt{r}-\sqrt{s}), \beta = (\sqrt{n}+\sqrt{m}+\sqrt{k}-\sqrt{r}-\sqrt{s})^2$. Let $\mathscr{D}_1$ denote the number of solutions of (2.3) with $|\alpha| \geqslant 2L^{-1/2}$, and $\mathscr{D}_2$ the number of solutions of (2.3) with $|\alpha| < 2L^{-1/2}$. By Lemma 2.4, we get

$$\begin{aligned}\mathscr{D}_1 &\ll (\Delta Q^{1/2}L + L^{1/2+\varepsilon})NMKRS \\ &\ll \Delta Q^{1/2}NMKLRS + NMKRSL^{1/2+\varepsilon}.\end{aligned}$$

Now we estimate $\mathscr{D}_2$. From $|\alpha| < 2L^{-1/2}$, we can get $K \asymp R$. If $\sqrt{n}+\sqrt{m}+\sqrt{k} = \sqrt{r}+\sqrt{s}$, then from (2.2), we get $\ell = q$. This contradicts to the fact that $|\sqrt{n}+\sqrt{m}+\sqrt{k}+\sqrt{\ell}-\sqrt{r}-\sqrt{s}\sqrt{q}| > 0$. Therefore, we have $\sqrt{n}+\sqrt{m}+\sqrt{k} \neq \sqrt{r}+\sqrt{s}$. By Lemma 2.1, we have $|\sqrt{n}+\sqrt{m}+\sqrt{k}-\sqrt{r}+\sqrt{s}| \gg S^{-15/2}$ for any such (n, m, k, r, s). By a splitting argument and Lemma 2.7, there exists a δ satisfying $S^{-15/2} \ll \delta \ll L^{-1/2}$, which holds

$$\begin{aligned}\mathscr{D}_2 &\ll \log 2S \cdot \sum_{\delta<|\sqrt{n}+\sqrt{m}+\sqrt{k}-\sqrt{r}+\sqrt{s}|\leqslant 2\delta} 1 \\ &\ll \log 2S \cdot (\delta S^{1/2}NMKR + NMKR) \\ &\ll L^{-1/2}S^{1/2+\varepsilon}NMKR + NMKRS^{\varepsilon} \\ &\ll NMKRS^{\varepsilon},\end{aligned}$$

which can be absorbed into the estimate of $\mathscr{D}_1$. This completes the proof of Lemma 2.8.

Lemma 2.9 *Suppose $1 \leqslant N \leqslant M \leqslant K \leqslant L, 1 \leqslant R \leqslant S \leqslant Q, L \asymp Q$ and $0 < \Delta \ll Q^{1/2}$. Let $\mathscr{A}_{2,\pm}(N, M, K, L, R, S, Q; \Delta)$ denote the number of solutions of the inequality*

$$0 < |\sqrt{n}+\sqrt{m}+\sqrt{k}+\sqrt{\ell}+\sqrt{r} \pm \sqrt{s}-\sqrt{q}| < \Delta$$

with $n \sim N, m \sim M, k \sim K, \ell \sim L, r \sim R, s \sim S, q \sim Q$. Then

$$\mathscr{A}_{2,\pm}(N, M, K, L, R, S, Q; \Delta) \ll \Delta Q^{1/2}NMKLRS + NMKRSL^{1/2+\varepsilon}.$$

In particular, if $\Delta Q^{1/2} \gg 1$, then

$$\mathscr{A}_{2,\pm}(N, M, K, L, R, S, Q; \Delta) \ll \Delta Q^{1/2}NMKLRS.$$

Proof. The proof of Lemma 2.9 is similar to that of Lemma 2.8, so we omit the details.

Lemma 2.10 *Suppose $N_j \geqslant 2 \quad (j = 1, 2, 3, 4, 5, 6, 7)$ are real numbers, $\Delta > 0$. let $\mathscr{A}_{\pm,\pm}(N_1.N_2, N_3, N_4, N_5, N_6, N_7; \Delta)$ denote the number of solutions of the inequality*

$$0 < |\sqrt{n_1}+\sqrt{n_2}+\sqrt{n_3}+\sqrt{n_4} \pm \sqrt{n_5} \pm \sqrt{n_6}-\sqrt{n_7}| < \Delta$$

with $n_j \sim N_j, (j = 1, 2, 3, 4, 5, 6, 7), n_j \in \mathbb{N}^*$. Then we have

$$\mathscr{A}_{\pm,\pm}(N_1.N_2, N_3, N_4, N_5, N_6, N_7; \Delta) \ll \prod_{j=1}^{7}\left(\Delta^{1/7}N_j^{13/14} + N_j^{5/7}\right)N_j^{\varepsilon}.$$

Proof. Taking $a = 6\Delta/5, \delta = \Delta/5$ in Lemma 2.5, there exists a function $\varphi_1(y)$, which is $\ell = [7\log(N_1N_2N_3N_4N_5N_6N_7)]$ times continuously differentiable such that

$$\begin{cases} \varphi_1(y) = 1, & \text{if} \quad |y| \leqslant \Delta, \\ 0 < \varphi_1(y) < 1, & \text{for} \quad \Delta < |y| < 7\Delta/5, \\ \varphi_1(y) = 0, & \text{for} \quad |y| \geqslant 7\Delta/5. \end{cases}$$

Let

$$\Phi_1(x) = \int_{-\infty}^{+\infty} e(-xy)\varphi_1(y)\mathrm{d}y,$$

then it satisfies

$$|\Phi_1(x)| \leqslant \min\left(\frac{12\Delta}{5}, \frac{1}{\pi|x|}, \frac{1}{\pi|x|}\left(\frac{5\ell}{2\pi|x|\Delta}\right)^{\ell}\right), \tag{2.4}$$

and

$$\varphi_1(y) = \int_{-\infty}^{+\infty} e(xy)\Phi_1(x)\mathrm{d}x. \tag{2.5}$$

Set

$$R_{\pm,\pm} = \sum_{\substack{n_j\sim N_j \\ j=1,2,3,4,5,6,7}} \varphi_1(\sqrt{n_1} + \sqrt{n_2} + \sqrt{n_3} + \sqrt{n_4} \pm \sqrt{n_5} \pm \sqrt{n_6} - \sqrt{n_7}).$$

By the definition of $\varphi_1(y)$, we get

$$\mathscr{A}_{\pm,\pm}(N_1.N_2, N_3, N_4, N_5, N_6, N_7; \Delta) \leqslant R_{\pm,\pm}. \tag{2.6}$$

We estimate $R_{-,-}$ first. By (2.5), we have

$$\begin{aligned} R_{-,-} &= \sum_{\substack{n_j\sim N_j \\ j=1,2,3,4,5,6,7}} \varphi_1(\sqrt{n_1} + \sqrt{n_2} + \sqrt{n_3} + \sqrt{n_4} - \sqrt{n_5} - \sqrt{n_6} - \sqrt{n_7}) \\ &= \sum_{\substack{n_j\sim N_j \\ j=1,2,3,4,5,6,7}} \int_{-\infty}^{+\infty} e\left(x(\sqrt{n_1} + \sqrt{n_2} + \sqrt{n_3} + \sqrt{n_4} - \sqrt{n_5} - \sqrt{n_6} - \sqrt{n_7})\right)\Phi_1(x)\mathrm{d}x. \end{aligned}$$

Let $S(x;N) := \sum_{n\sim N} e(x\sqrt{n})$, we have

$$R_{-,-} = \int_{-\infty}^{+\infty} S(x;N_1)S(x;N_2)S(x;N_3)S(x;N_4)\overline{S(x;N_5)S(x;N_6)S(x;N_7)}\Phi_1(x)\mathrm{d}x,$$

if we notice that $\overline{S(x;N)} = \sum_{n\sim N} e(-x\sqrt{n})$. Applying Hölders inequality, we get

$$R_{-,-} \leqslant \prod_{j=1}^{7}\left(\int_{-\infty}^{+\infty} |S(x;N_j)|^7|\Phi_1(x)|\mathrm{d}x\right)^{1/7}. \tag{2.7}$$

Let

$$T(N) := \int_{0}^{+\infty} |S(x;N)|^7|\Phi_1(x)|\mathrm{d}x.$$

It is sufficient to estimate $T(N)$, where $N = N_j$ for some $j \in \{1,2,3,4,5,6,7\}$. Let

$$K := \frac{100\ell^2}{\Delta}, \quad \ell = [7\log(N_1N_2N_3N_4N_5N_6N_7)]; \quad K_0 := N^{1/2}.$$

Using the trivial estimate $S(x;N) \ll N$ and the estimate

$$|\Phi_1(x)| \leqslant \frac{1}{\pi|x|}\left(\frac{5\ell}{2\pi|x|\Delta}\right)^{\ell},$$

we have

$$\begin{aligned}\int_K^{\infty} |S(x;N)|^7|\Phi_1(x)|\mathrm{d}x &\ll N^7\int_K^{\infty}|\Phi_1(x)|\mathrm{d}x \ll N^7\left(\frac{5\ell}{2\pi\Delta}\right)^{\ell}\int_K^{\infty}\frac{1}{x^{\ell+1}}\mathrm{d}x\\ &\ll \frac{N^7 5^{\ell}}{\ell}\left(\frac{\ell}{2\pi K\Delta}\right)^{\ell} \ll \frac{N^7 5^{\ell}}{\ell}\frac{1}{\ell^{\ell}}\\ &\ll \frac{N^7(N_1\cdots N_7)^{7\log 5}}{(N_1\cdots N_7)^{7\log 7}(N_1\cdots N_7)^{7\log\log(N_1\cdots N_7)}} \ll 1. \end{aligned} \tag{2.8}$$

For the mean square of $S(x;N)$, we have

$$\begin{aligned}\int_0^{K_0}|S(x;N)|^2\mathrm{d}x &= \int_0^{K_0}\sum_{n\sim N}\sum_{m\sim N} e(x(\sqrt{n}-\sqrt{m}))\mathrm{d}x\\ &= \sum_{n\sim N}\sum_{m\sim N}\int_0^{K_0} e(x(\sqrt{n}-\sqrt{m}))\mathrm{d}x\\ &= \Big\{\sum_{\substack{n\sim N\\ n=m}}\sum_{m\sim N} + \sum_{\substack{n\sim N\\ n\neq m}}\sum_{m\sim N}\Big\}\int_0^{K_0} e(x(\sqrt{n}-\sqrt{m}))\mathrm{d}x.\end{aligned}$$

If we use the trivial estimate $S(x;N) \ll N$, then

$$\sum_{\substack{n\sim N\\ n=m}}\sum_{m\sim N}\int_0^{K_0} e(x(\sqrt{n}-\sqrt{m}))\mathrm{d}x \ll N^{3/2}.$$

For the case $n \neq m$, we have

$$\begin{aligned}&\sum_{\substack{n\sim N\\ n\neq m}}\sum_{m\sim N}\int_0^{K_0} e(x(\sqrt{n}-\sqrt{m}))\mathrm{d}x\\ \ll &\sum_{\substack{n\sim N\\ n\neq m}}\sum_{m\sim N}\frac{1}{|\sqrt{n}-\sqrt{m}|} \ll N^{1/2}\sum_{\substack{n\sim N\\ n\neq m}}\sum_{m\sim N}\frac{1}{|n-m|}\\ \ll &\ N^{1/2}\sum_{n\sim N}\sum_{1\leqslant r\ll N}\frac{1}{r} \ll N^{3/2}\log N.\end{aligned}$$

Therefore, we have

$$\int_0^{K_0}|S(x;N)|^2\mathrm{d}x \ll N^{3/2}\log N. \tag{2.9}$$

If we notice $|\Phi_1(x)| \ll \Delta$ from (2.4) and the trivial estimate $S(x;N) \ll N$, then

$$\begin{aligned}
&\int_0^{K_0} |S(x;N)|^7 |\Phi_1(x)| \mathrm{d}x \\
\ll\ & \Delta N^5 \int_0^{K_0} |S(x;N)|^2 \mathrm{d}x \\
\ll\ & \Delta N^{13/2} \log N. \qquad (2.10)
\end{aligned}$$

If $K \leqslant K_0$, then from (2.8) and (2.10) we get

$$T(N) \ll \Delta N^{13/2} \log N. \qquad (2.11)$$

Now suppose $K_0 < K$. By a splitting argument, we have

$$\int_{K_0}^{K} |S(x;N)|^7 |\Phi_1(x)| \mathrm{d}x \ll \Delta \log K \times \max_{K_0 \leqslant U \leqslant K} \int_U^{2U} |S(x;N)|^7 \mathrm{d}x. \qquad (2.12)$$

From Lemma 2.1, we can get $\Delta^{-1} \ll \max(N_1, N_2, N_3, N_4, N_5, N_6, N_7)^{63/2}$ and thus $\log K \ll \ell^2$. On the other hand, we have

$$\begin{aligned}
\int_U^{2U} |S(x;N)|^7 \mathrm{d}x &\ll \max_{U \leqslant x \leqslant 2U} |S(x;N)|^2 \times \int_U^{2U} |S(x;N)|^5 \mathrm{d}x \\
&\ll N^2 (N^{9/2} + UN^3) N^{\varepsilon} \\
&\ll (N^{13/2} + UN^5) N^{\varepsilon}, \qquad (2.13)
\end{aligned}$$

which is derived from equation (2.18) of Zhang and Zhai [21].

From (2.12) and (2.13) and noticing $\Delta K \ll \ell^2$, we get

$$\begin{aligned}
\int_{K_0}^{K} |S(x;N)|^7 |\Phi_1(x)| \mathrm{d}x &\ll \Delta \log K \times (N^{13/2} + UN^5) N^{\varepsilon} \\
&\ll (\Delta N^{13/2} + N^5) N^{\varepsilon},
\end{aligned}$$

which combining (2.8) and (2.10) gives

$$T(N) = \int_0^{\infty} |S(x;N)|^7 |\Phi_1(x)| \mathrm{d}x \ll (\Delta N^{13/2} + N^5) N^{\varepsilon}. \qquad (2.14)$$

From (2.6), (2.7) and (2.14), we get the result of Lemma 2.10 for the case "$-,-$". By noting the properties of conjugation, the estimates of other cases are exactly the same as that of the case "$-,-$". This completes the proof of Lemma 2.10.

3 Proof of Theorem

In this section, we shall prove the theorem. We begin with the following truncated form of the Voronoï's formula ([7], equation (2.25)), i.e.

$$\Delta(x) = \frac{1}{\sqrt{2}\pi} \sum_{n \leqslant N} \frac{d(n)}{n^{3/4}} x^{1/4} \cos(4\pi\sqrt{nx} - \pi/4) + O(x^{1/2+\varepsilon} N^{-1/2}), \qquad (3.1)$$

for $1 \leqslant N \ll x$. Set $\Delta(x) := R_1 + R_2$, where

$$R_1 := R_1(x) = \frac{1}{\sqrt{2}\pi} \sum_{n \leqslant y} \frac{d(n)}{n^{3/4}} x^{1/4} \cos(4\pi\sqrt{nx} - \pi/4), \quad R_2 := R_2(x) = \Delta(x) - R_1.$$

Take $y = T^{1/4}$. By the elementary estimate $(a+b)^7 - a^7 \ll |b|a^6 + |b|^7$, we have

$$\int_1^T \Delta^7(x)\mathrm{d}x = \int_1^T R_1^7(x)\mathrm{d}x + O\left(\int_1^T |R_1|^6|R_2|\mathrm{d}x + \int_1^T |R_2|^7\mathrm{d}x\right).$$

By a splitting argument, it is sufficient to prove the result in the interval $[T, 2T]$. We will divide the process of the proof of the theorem into two parts.

Proposition 3.1 *For fixed $T \geqslant 10, N = T, y = T^{1/4}$, we have*

$$\int_T^{2T} R_1^7 \mathrm{d}x = \frac{7(5s_{7;3}(d) - 3s_{7;2}(d) - s_{7;1}(d))}{2816\pi^7} T^{11/4} + O(T^{11/4 - 1/336 + \varepsilon}). \tag{3.2}$$

Proof. Let

$$g := g(n,m,k,\ell,r,s,q) = \begin{cases} \frac{d(n)d(m)d(k)d(\ell)d(r)d(s)d(q)}{(nmk\ell rsq)^{3/4}}, & \text{if} \quad n,m,k,\ell,r,s,q \leqslant y, \\ 0, & \text{otherwise.} \end{cases}$$

According to the elementary formula

$$\cos a_1 \cos a_2 \cdots \cos a_h = \frac{1}{2^{h-1}} \sum_{(i_1,i_2,\cdots,i_{h-1}) \in \{0,1\}^{h-1}} \cos\left(a_1 + (-1)^{i_1} a_2 + \cdots + (-1)^{i_{h-1}} a_h\right),$$

we can write

$$R_1^7 = S_1(x) + S_2(x) + S_3(x) + S_4(x) + S_5(x) + S_6(x) + S_7(x),$$

where

$$\begin{aligned}
S_1(x) &:= \frac{35}{64}\cos\frac{\pi}{4} \sum_{\substack{n,m,k,\ell,r,s,q \leqslant y \\ \sqrt{n}+\sqrt{m}+\sqrt{k}+\sqrt{\ell} = \sqrt{r}+\sqrt{s}+\sqrt{q}}} g x^{7/4}, \\
S_2(x) &:= \frac{35}{64} \sum_{\substack{n,m,k,\ell,r,s,q \leqslant y \\ \sqrt{n}+\sqrt{m}+\sqrt{k}+\sqrt{\ell} \neq \sqrt{r}+\sqrt{s}+\sqrt{q}}} g x^{7/4} \\
&\quad \times \cos\left(4\pi\left(\sqrt{n}+\sqrt{m}+\sqrt{k}+\sqrt{\ell}-\sqrt{r}-\sqrt{s}-\sqrt{q}\right)\sqrt{x} - \frac{\pi}{4}\right), \\
S_3(x) &:= \frac{21}{64}\cos\frac{3\pi}{4} \sum_{\substack{n,m,k,\ell,r,s,q \leqslant y \\ \sqrt{n}+\sqrt{m}+\sqrt{k}+\sqrt{\ell}+\sqrt{r} = \sqrt{s}+\sqrt{q}}} g x^{7/4}, \\
S_4(x) &:= \frac{21}{64} \sum_{\substack{n,m,k,\ell,r,s,q \leqslant y \\ \sqrt{n}+\sqrt{m}+\sqrt{k}+\sqrt{\ell}+\sqrt{r} \neq \sqrt{s}+\sqrt{q}}} g x^{7/4} \\
&\quad \times \cos\left(4\pi\left(\sqrt{n}+\sqrt{m}+\sqrt{k}+\sqrt{\ell}+\sqrt{r}-\sqrt{s}-\sqrt{q}\right)\sqrt{x} - \frac{3\pi}{4}\right),
\end{aligned}$$

$$
\begin{aligned}
S_5(x) &:= \frac{7}{64}\cos\frac{5\pi}{4} \sum_{\substack{n,m,k,\ell,r,s,q\leqslant y\\ \sqrt{n}+\sqrt{m}+\sqrt{k}+\sqrt{\ell}+\sqrt{r}+\sqrt{s}=\sqrt{q}}} g x^{7/4},\\
S_6(x) &:= \frac{7}{64} \sum_{\substack{n,m,k,\ell,r,s,q\leqslant y\\ \sqrt{n}+\sqrt{m}+\sqrt{k}+\sqrt{\ell}+\sqrt{r}+\sqrt{s}\neq\sqrt{q}}} g x^{7/4}\\
&\quad \times \cos\left(4\pi\left(\sqrt{n}+\sqrt{m}+\sqrt{k}+\sqrt{\ell}+\sqrt{r}+\sqrt{s}-\sqrt{q}\right)\sqrt{x}-\frac{5\pi}{4}\right),\\
S_7(x) &:= \frac{1}{64} \sum_{n,m,k,\ell,r,s,q\leqslant y} g x^{7/4}\\
&\quad \times \cos\left(4\pi\left(\sqrt{n}+\sqrt{m}+\sqrt{k}+\sqrt{\ell}+\sqrt{r}+\sqrt{s}+\sqrt{q}\right)\sqrt{x}-\frac{7\pi}{4}\right).
\end{aligned}
$$

By Lemma 2.6, we get

$$
\begin{aligned}
\int_T^{2T} S_1(x)\mathrm{d}x &= \frac{35\sqrt{2}}{128} s_{7;3}(d;y)\int_T^{2T} x^{7/4}\mathrm{d}x\\
&= \frac{35\sqrt{2}}{128} s_{7;3}(d)\int_T^{2T} x^{7/4}\mathrm{d}x + O\left(T^{11/4}y^{-1/2+\varepsilon}\right)\\
&= \frac{35\sqrt{2}}{128} s_{7;3}(d)\int_T^{2T} x^{7/4}\mathrm{d}x + O\left(T^{11/4-1/8+\varepsilon}\right). \qquad (3.3)
\end{aligned}
$$

Similarly, we can get

$$
\int_T^{2T} S_3(x)\mathrm{d}x = -\frac{21\sqrt{2}}{128} s_{7;2}(d)\int_T^{2T} x^{7/4}\mathrm{d}x + O\left(T^{11/4-1/8+\varepsilon}\right) \qquad (3.4)
$$

and

$$
\int_T^{2T} S_5(x)\mathrm{d}x = -\frac{7\sqrt{2}}{128} s_{7;1}(d)\int_T^{2T} x^{7/4}\mathrm{d}x + O\left(T^{11/4-1/8+\varepsilon}\right). \qquad (3.5)
$$

We now proceed to consider the contribution of $S_7(x)$. Applying Lemma 2.3, we have

$$
\begin{aligned}
\int_T^{2T} S_7(x)\mathrm{d}x &\ll \sum_{n,m,k,\ell,r,s,q\leqslant y} \frac{gT^{9/4}}{\sqrt{n}+\sqrt{m}+\sqrt{k}+\sqrt{\ell}+\sqrt{r}+\sqrt{s}+\sqrt{q}}\\
&\ll T^{9/4+\varepsilon} \sum_{n\leqslant m\leqslant k\leqslant \ell\leqslant r\leqslant s\leqslant q\leqslant y} \frac{1}{(nmk\ell rsq)^{3/4}q^{1/2}}\\
&\ll T^{9/4+\varepsilon}y^{5/4} \ll T^{41/16+\varepsilon}. \qquad (3.6)
\end{aligned}
$$

Now we consider the contribution of $S_2(x)$. By the first derivative test, we get

$$
\begin{aligned}
\int_T^{2T} S_2(x)\mathrm{d}x &\ll \sum_{\substack{n,m,k,\ell,r,s,q\leqslant y\\ \sqrt{n}+\sqrt{m}+\sqrt{k}+\sqrt{\ell}\neq\sqrt{r}+\sqrt{s}+\sqrt{q}}} g\\
&\quad \times \min\left(T^{11/4}, \frac{T^{9/4}}{|\sqrt{n}+\sqrt{m}+\sqrt{k}+\sqrt{\ell}-\sqrt{r}-\sqrt{s}-\sqrt{q}|}\right)\\
&\ll x^{\varepsilon}G(N,M,K,L,R,S,Q), \qquad (3.7)
\end{aligned}
$$

where

$$G(N,M,K,L,R,S,Q) = \sum_{\substack{\sqrt{n}+\sqrt{m}+\sqrt{k}+\sqrt{\ell}\neq\sqrt{r}+\sqrt{s}+\sqrt{q}\\ n\sim N,m\sim M,k\sim K,\ell\sim L,r\sim R,s\sim S,q\sim Q\\ 1\leqslant N\leqslant M\leqslant K\leqslant L\leqslant y\\ 1\leqslant R\leqslant S\leqslant Q\leqslant y}} g \times \min\left(T^{11/4}, \frac{T^{9/4}}{|\sqrt{n}+\sqrt{m}+\sqrt{k}+\sqrt{\ell}-\sqrt{r}-\sqrt{s}-\sqrt{q}|}\right).$$

If $L \geqslant 100Q$, then $|\sqrt{n}+\sqrt{m}+\sqrt{k}+\sqrt{\ell}-\sqrt{r}-\sqrt{s}-\sqrt{q}| \gg L^{1/2}$, so the trivial estimate yields

$$G(N,M,K,L,R,S,Q) \ll \frac{T^{9/4+\varepsilon}NMKLRSQ}{(NMKLRSQ)^{3/4}L^{1/2}} \ll T^{9/4+\varepsilon}y^{5/4} \ll T^{41/16+\varepsilon}.$$

If $Q \geqslant 100L$, we can get the same estimate. So later we always suppose that $L \asymp Q$. Let $\eta = \sqrt{n}+\sqrt{m}+\sqrt{k}+\sqrt{\ell}-\sqrt{r}-\sqrt{s}-\sqrt{q}$. Write

$$G(N,M,K,L,R,S,Q) = G_1 + G_2 + G_3, \tag{3.8}$$

where

$$\begin{aligned} G_1 &:= T^{11/4} \sum_{0<|\eta|\leqslant T^{-1/2}} g, \\ G_2 &:= T^{9/4} \sum_{T^{-1/2}<|\eta|\leqslant 1} g|\eta|^{-1}, \\ G_3 &:= T^{9/4} \sum_{|\eta|>1} g|\eta|^{-1}. \end{aligned}$$

We estimate G_1 first. From $|\eta| \leqslant T^{-1/2}$, we get $Q \gg T^{1/63}$ via Lemma 2.1. By Lemma 2.8, we get

$$\begin{aligned} G_1 &\ll \frac{T^{11/4+\varepsilon}}{(NMKLRSQ)^{3/4}}\mathscr{A}_1(N,M,K,L,R,S,Q;T^{-1/2}) \\ &\ll \frac{T^{11/4+\varepsilon}}{(NMKLRSQ)^{3/4}}\left(T^{-1/2}Q^{1/2}NMKLRS + NMKRSL^{1/2}\right) \\ &\ll T^{9/4+\varepsilon}y^{5/4} + T^{11/4+\varepsilon}(NMKRS)^{1/4}L^{-1} \\ &\ll T^{41/16+\varepsilon} + T^{11/4+\varepsilon}(NMKRS)^{1/4}L^{-1}. \end{aligned} \tag{3.9}$$

By Lemma 2.10, we get

$$\begin{aligned} G_1 &\ll \frac{T^{11/4+\varepsilon}}{(NMKLRSQ)^{3/4}}\mathscr{A}_{-,-}(N,M,K,L,R,S,Q;T^{-1/2}) \\ &\ll \frac{T^{11/4+\varepsilon}}{(NMKLRSQ)^{3/4}}(T^{-1/14}N^{13/14}+N^{5/7})(T^{-1/14}M^{13/14}+M^{5/7}) \\ &\quad \times(T^{-1/14}K^{13/14}+K^{5/7})(T^{-1/14}R^{13/14}+R^{5/7})(T^{-1/14}S^{13/14}+S^{5/7})(T^{-1/7}Q^{13/7}+Q^{10/7}) \\ &\ll T^{11/4+\varepsilon}Q^{-1/14}(NMKRS)^{-1/28}(T^{-1/7}Q^{3/7}+1)(T^{-1/14}N^{3/14}+1) \\ &\quad \times(T^{-1/14}M^{3/14}+1)(T^{-1/14}K^{3/14}+1)(T^{-1/14}R^{3/14}+1)(T^{-1/14}S^{3/14}+1) \\ &\ll T^{11/4+\varepsilon}Q^{-1/14}(NMKRS)^{-1/28} \end{aligned} \tag{3.10}$$

by noting that $T^{-1/7}D^{3/7} \ll 1$ for $D = Q, N, M, K, R, S$. From (3.9) and (3.10), we get

$$\begin{aligned} G_1 &\ll T^{41/16+\varepsilon} + T^{11/4+\varepsilon} \cdot \min\left(\frac{(NMKRS)^{1/4}}{L}, \frac{1}{Q^{1/14}(NMKRS)^{1/28}}\right) \\ &\ll T^{41/16+\varepsilon} + T^{11/4+\varepsilon}\left(\frac{(NMKRS)^{1/4}}{L}\right)^{1/8}\left(\frac{1}{Q^{1/14}(NMKRS)^{1/28}}\right)^{7/8} \\ &\ll T^{41/16+\varepsilon} + T^{11/4+\varepsilon}Q^{-3/16} \\ &\ll T^{11/4-1/336+\varepsilon}, \end{aligned} \tag{3.11}$$

if we notice the fact that $Q \gg T^{1/63}$.

Now we estimate G_2. We also suppose $K \leqslant R$ and the other cases are the same. By a splitting argument, we get the estimate

$$G_2 \ll \frac{T^{9/4+\varepsilon}}{(NMKLRSQ)^{3/4}\delta} \times \sum_{\substack{\delta<|\eta|\leqslant 2\delta \\ |\eta|\neq 0}} 1$$

for some $T^{-1/2} \leqslant \delta \leqslant 1$. By Lemma 2.8, we get

$$\begin{aligned} G_2 &\ll \frac{T^{9/4+\varepsilon}}{(NMKLRSQ)^{3/4}\delta}\mathscr{A}_1(N, M, K, L, R, S, Q; 2\delta) \\ &\ll \frac{T^{9/4+\varepsilon}}{(NMKLRSQ)^{3/4}\delta}\left(\delta Q^{1/2}NMKLRS + NMKRSL^{1/2}\right) \\ &\ll T^{9/4+\varepsilon}y^{5/4} + T^{9/4+\varepsilon}(NMKRS)^{1/4}(L\delta)^{-1} \\ &\ll T^{41/16+\varepsilon} + T^{9/4+\varepsilon}(NMKRS)^{1/4}(L\delta)^{-1}. \end{aligned} \tag{3.12}$$

On the other hand, by Lemma 2.10, we have

$$\begin{aligned} G_2 &\ll \frac{T^{9/4+\varepsilon}}{(NMKLRSQ)^{3/4}\delta}\mathscr{A}_{-,-}(N, M, K, L, R, S, Q; 2\delta) \\ &\ll \frac{T^{9/4+\varepsilon}}{(NMKLRSQ)^{3/4}\delta}(\delta^{1/7}N^{13/14} + N^{5/7})(\delta^{1/7}M^{13/14} + M^{5/7}) \\ &\quad \times(\delta^{1/7}K^{13/14} + K^{5/7})(\delta^{1/7}R^{13/14} + R^{5/7})(\delta^{1/7}S^{13/14} + S^{5/7})(\delta^{2/7}Q^{13/7} + Q^{10/7}) \\ &\ll T^{9/4+\varepsilon}\delta^{-1}(\delta^{1/7}N^{5/28} + N^{-1/28})(\delta^{1/7}M^{5/28} + M^{-1/28})(\delta^{1/7}K^{5/28} + K^{-1/28}) \\ &\quad \times(\delta^{1/7}R^{5/28} + R^{-1/28})(\delta^{1/7}S^{5/28} + S^{-1/28})(\delta^{2/7}Q^{5/14} + Q^{-1/14}) \\ &\ll T^{9/4+\varepsilon}\delta^{-1}Q^{-1/14}(NMKRS)^{-1/28}(\delta^{2/7}Q^{3/7} + 1)(\delta^{1/7}N^{3/14} + 1) \\ &\quad \times(\delta^{1/7}M^{3/14} + 1)(\delta^{1/7}K^{3/14} + 1)(\delta^{1/7}R^{3/14} + 1)(\delta^{1/7}S^{3/14} + 1) \\ &\ll T^{9/4+\varepsilon}\delta^{-1}Q^{-1/14}(NMKRS)^{-1/28}(\delta^{2/7}Q^{3/7} + 1)(\delta^{5/7}(NMKRS)^{3/14} \\ &\quad +\delta^{4/7}(MKRS)^{3/14} + \delta^{3/7}(KRS)^{3/14} + \delta^{2/7}(RS)^{3/14} + \delta^{1/7}(S)^{3/14} + 1) \\ &\ll T^{9/4+\varepsilon}\delta^{-2/7}Q^{-1/14}(NMKRS)^{5/28}(\delta^{2/7}Q^{3/7} + 1) \\ &\quad +T^{9/4+\varepsilon}\delta^{-1}Q^{-1/14}(NMKRS)^{-1/28}(\delta^{2/7}Q^{3/7} + 1) \\ &\quad \times(\delta^{4/7}(MKRS)^{3/14} + \delta^{3/7}(KRS)^{3/14} + \delta^{2/7}(RS)^{3/14} + \delta^{1/7}S^{3/14} + 1) \end{aligned}$$

$$\begin{aligned}
&\ll T^{9/4+\varepsilon}y^{5/4}+T^{9/4+\varepsilon}T^{1/7}y^{23/28}+T^{9/4+\varepsilon}\delta^{-1}Q^{-1/14}(NMKRS)^{-1/28}(\delta^{2/7}Q^{3/7}+1)\\
&\quad\times(\delta^{4/7}Q^{6/7}+\delta^{3/7}Q^{9/14}+\delta^{2/7}Q^{3/7}+\delta^{1/7}Q^{3/14}+1)\\
&\ll T^{291/112+\varepsilon}+T^{9/4+\varepsilon}\delta^{-1}Q^{-1/14}(NMKRS)^{-1/28}(\delta^{6/7}Q^{9/7}+1). \qquad (3.13)
\end{aligned}$$

From (3.12) and (3.13), we get

$$\begin{aligned}
G_2 &\ll T^{291/112+\varepsilon}+T^{9/4+\varepsilon}\delta^{-1}\cdot\min\left(\frac{(NMKRS)^{1/4}}{L},\frac{\delta^{6/7}Q^{9/7}+1}{Q^{1/14}(NMKRS)^{1/28}}\right)\\
&\ll T^{291/112+\varepsilon}+T^{9/4+\varepsilon}\delta^{-1}\left(\frac{(NMKRS)^{1/4}}{L}\right)^{1/8}\left(\frac{\delta^{6/7}Q^{9/7}+1}{Q^{1/14}(NMKRS)^{1/28}}\right)^{7/8}\\
&\ll T^{291/112+\varepsilon}+T^{9/4+\varepsilon}\delta^{-1}Q^{-3/16}(\delta^{3/4}Q^{9/8}+1). \qquad (3.14)
\end{aligned}$$

If $\delta\gg Q^{-3/2}$, then $\delta^{3/4}Q^{9/8}\gg 1$ and (3.14) implies (recall $\delta\gg T^{-1/2}$)

$$\begin{aligned}
G_2 &\ll T^{291/112+\varepsilon}+T^{9/4+\varepsilon}\delta^{-1/4}Q^{15/16}\\
&\ll T^{291/112+\varepsilon}+T^{9/4+1/8+\varepsilon}y^{15/16}\\
&\ll T^{167/64+\varepsilon}. \qquad (3.15)
\end{aligned}$$

If $\delta\ll Q^{-3/2}$, then $\delta^{3/4}Q^{9/8}\ll 1$ and (3.14) implies

$$G_2\ll T^{291/112+\varepsilon}+T^{9/4+\varepsilon}\delta^{-1}Q^{-3/16}. \qquad (3.16)$$

By Lemma 2.1, we have $2\delta\geqslant|\eta|\gg Q^{-63/2}$, which implies $\delta^{-1}\ll Q^{63/2}$. Therefore, we get

$$\delta^{-1}\ll\min\left(Q^{63/2},T^{1/2}\right).$$

From the above estimate and (3.16), we get

$$\begin{aligned}
G_2 &\ll T^{291/112+\varepsilon}+\min\left(T^{9/4+\varepsilon}Q^{501/16},T^{11/4+\varepsilon}Q^{-3/16}\right)\\
&\ll T^{291/112+\varepsilon}+\left(T^{9/4+\varepsilon}Q^{501/16}\right)^{1/168}\left(T^{11/4+\varepsilon}Q^{-3/16}\right)^{167/168}\\
&\ll T^{11/4-1/336+\varepsilon}. \qquad (3.17)
\end{aligned}$$

For G_3, by a splitting argument and Lemma 2.8 (notice $|\eta|\gg 1$),we get

$$\begin{aligned}
G_3 &\ll \frac{T^{9/4+\varepsilon}}{(NMKLRSQ)^{3/4}\delta}\times\sum_{\substack{\delta<|\eta|\leqslant 2\delta\\ |\delta|\gg 1}}1\\
&\ll \frac{T^{9/4+\varepsilon}}{(NMKLRSQ)^{3/4}}Q^{1/2}NMKLRS\\
&\ll T^{9/4+\varepsilon}y^{5/4}\\
&\ll T^{41/16+\varepsilon}. \qquad (3.18)
\end{aligned}$$

Combining (3.7), (3.8), (3.11), (3.17) and (3.18), we get

$$\int_T^{2T}S_2(x)\mathrm{d}x\ll T^{11/4-1/336+\varepsilon}. \qquad (3.19)$$

In the same way, we can prove that

$$\int_T^{2T} S_4(x)\mathrm{d}x \ll T^{11/4-1/336+\varepsilon} \tag{3.20}$$

and

$$\int_T^{2T} S_6(x)\mathrm{d}x \ll T^{11/4-1/336+\varepsilon}, \tag{3.21}$$

if we use Lemma 2.9 instead of Lemma 2.8.

Now Proposition 3.1, i.e. equation (3.2), follows from (3.3)-(3.6) and (3.19)-(3.21).

Proposition 3.2 *For $T^\varepsilon \ll y \ll T^{1/4}$, we have*

$$\int_T^{2T} |R_2|^7\mathrm{d}x \ll T^{11/4-1/336+\varepsilon} \tag{3.22}$$

and

$$\int_T^{2T} |R_1|^6|R_2|\mathrm{d}x \ll T^{11/4-1/336+\varepsilon}. \tag{3.23}$$

Proof. First, for $T \leqslant x \leqslant 2T$, we have

$$\begin{aligned} R_2 &= \frac{1}{\sqrt{2}\pi}\sum_{y<n\leqslant T}\frac{d(n)}{n^{3/4}}x^{1/4}\cos(4\pi\sqrt{nx}-\pi/4)+O(x^{1/2+\varepsilon}T^{-1/2}) \\ &\ll x^{1/4}\left|\sum_{y<n\leqslant T}\frac{d(n)}{n^{3/4}}\cos(4\pi\sqrt{nx}-\pi/4)\right|+T^\varepsilon \\ &\ll x^{1/4}\left|\sum_{y<n\leqslant T}\frac{d(n)}{n^{3/4}}e\left(2\sqrt{nx}\right)\right|+T^\varepsilon. \end{aligned}$$

Therefore, one has

$$\begin{aligned} \int_T^{2T}|R_2|^2\mathrm{d}x &\ll T^{1+\varepsilon}+\int_T^{2T}x^{1/2}\left|\sum_{y<n\leqslant T}\frac{d(n)}{n^{3/4}}e\left(2\sqrt{nx}\right)\right|^2\mathrm{d}x \\ &\ll T^{1+\varepsilon}+T^{1/2}\int_T^{2T}\sum_{y<n\leqslant T}\sum_{y<m\leqslant T}\frac{d(n)d(m)}{(nm)^{3/4}}e\left(2(\sqrt{n}-\sqrt{m})\sqrt{x}\right)\mathrm{d}x \\ &= T^{1+\varepsilon}+T^{1/2}\sum_{y<n\leqslant T}\sum_{y<m\leqslant T}\frac{d(n)d(m)}{(nm)^{3/4}}\int_T^{2T}e\left(2(\sqrt{n}-\sqrt{m})\sqrt{x}\right)\mathrm{d}x \\ &\ll T^{1+\varepsilon}+T^{3/2}\sum_{y<n\leqslant T}\frac{d^2(n)}{n^{3/2}}+T\sum_{\substack{y<n\leqslant T\\ n\neq m}}\sum_{y<m\leqslant T}\frac{d(n)d(m)}{(nm)^{3/4}|\sqrt{n}-\sqrt{m}|} \\ &\ll T^{1+\varepsilon}+T^{3/2}\sum_{n>y}\frac{d^2(n)}{n^{3/2}}+T\sum_{m<n\leqslant T}\frac{d(n)d(m)}{(nm)^{3/4}|\sqrt{n}-\sqrt{m}|} \\ &\ll T^{1+\varepsilon}+T^{3/2}y^{-1/2}\log^3 y+T\sum_{m<n\leqslant T}\frac{d(n)d(m)}{(nm)^{3/4}(\sqrt{n}-\sqrt{m})} \\ &\ll T^{3/2}y^{-1/2}\log^3 T+T\sum_{m<n\leqslant T}\frac{d(n)d(m)}{(nm)^{3/4}(\sqrt{n}-\sqrt{m})}. \end{aligned} \tag{3.24}$$

For the last sum in (3.24), we have

$$\sum_{m<n\leqslant T}\frac{d(n)d(m)}{(nm)^{3/4}(\sqrt{n}-\sqrt{m})}:=\Sigma_1+\Sigma_2, \tag{3.25}$$

where

$$\Sigma_1:=\sum_{\substack{m<n\leqslant T\\ \sqrt{n}-\sqrt{m}>\frac{1}{10}(nm)^{1/4}}}\frac{d(n)d(m)}{(nm)^{3/4}(\sqrt{n}-\sqrt{m})},$$

$$\Sigma_2:=\sum_{\substack{m<n\leqslant T\\ 0<\sqrt{n}-\sqrt{m}\leqslant\frac{1}{10}(nm)^{1/4}}}\frac{d(n)d(m)}{(nm)^{3/4}(\sqrt{n}-\sqrt{m})}.$$

For Σ_1, one has

$$\Sigma_1\ll\sum_{n\leqslant T}\sum_{m<n}\frac{d(n)d(m)}{nm}\ll\sum_{n\leqslant T}\frac{d(n)}{n}\sum_{m<n}\frac{d(m)}{m}\ll\log^4 T. \tag{3.26}$$

For Σ_2, we can write

$$\Sigma_2\ll\sum_{\substack{m<n\leqslant T\\ 0<\sqrt{n}-\sqrt{m}\leqslant\frac{1}{10}(nm)^{1/4}}}\frac{d^2(n)+d^2(m)}{(nm)^{3/4}(\sqrt{n}-\sqrt{m})}:=\Sigma_{21}+\Sigma_{22}, \tag{3.27}$$

where

$$\Sigma_{21}:=\sum_{\substack{m<n\leqslant T\\ 0<\sqrt{n}-\sqrt{m}\leqslant\frac{1}{10}(nm)^{1/4}}}\frac{d^2(n)}{(nm)^{3/4}(\sqrt{n}-\sqrt{m})},$$

$$\Sigma_{22}:=\sum_{\substack{m<n\leqslant T\\ 0<\sqrt{n}-\sqrt{m}\leqslant\frac{1}{10}(nm)^{1/4}}}\frac{d^2(m)}{(nm)^{3/4}(\sqrt{n}-\sqrt{m})}.$$

By the mean value theorem and taking $n-m=r$, we get

$$\Sigma_{21}\ll\sum_{n\leqslant T}\sum_{m<n}\frac{d^2(n)}{n(n-m)}\ll\sum_{n\leqslant T}\frac{d^2(n)}{n}\sum_{1\leqslant r\ll T}\frac{1}{r}\ll\log^5 T, \tag{3.28}$$

$$\Sigma_{22}\ll\sum_{n\leqslant T}\sum_{m<n}\frac{d^2(m)}{m(n-m)}\ll\sum_{m<T}\frac{d^2(m)}{m}\sum_{1\leqslant r\ll T}\frac{1}{r}\ll\log^5 T. \tag{3.29}$$

By the estimate (3.24)-(3.29), we get

$$\int_T^{2T}|R_2|^2\mathrm{d}x\ll T^{3/2}y^{-1/2}\log^3 T. \tag{3.30}$$

Taking $y=T^{1/4}$, we get $R_1\ll T^{1/4}y^{1/4}\ll T^{131/416}$, and therefore $R_2\ll T^{131/416+\varepsilon}$. Using Ivić's large-value technique directly to R_2 without modifications, one can get that the estimate

$$\int_T^{2T}|R_2|^{A_0}\mathrm{d}x\ll T^{1+\frac{A_0}{4}+\varepsilon} \tag{3.31}$$

holds with $A_0 = \frac{184}{19}, T^{\varepsilon} \ll y \ll T^{1/4}$. Then, for $2 < A < A_0$, by (3.30), (3.31) and Hölder's inequality, we have

$$\begin{aligned}\int_T^{2T} |R_2|^A \mathrm{d}x &= \int_T^{2T} |R_2|^{\frac{2(A_0-A)}{A_0-2}+\frac{A_0(A-2)}{A_0-2}} \mathrm{d}x \\ &\ll \left(\int_T^{2T} |R_2|^2 \mathrm{d}x\right)^{\frac{A_0-A}{A_0-2}} \left(\int_T^{2T} |R_2|^{A_0} \mathrm{d}x\right)^{\frac{A-2}{A_0-2}} \\ &\ll T^{1+\frac{A}{4}+\varepsilon} y^{-\frac{A_0-A}{2(A_0-2)}}.\end{aligned}$$

Especially, we take $A = 7, y = T^{1/4}$ and get

$$\begin{aligned}\int_T^{2T} |R_2|^7 \mathrm{d}x &\ll T^{11/4+\varepsilon} y^{-51/292} \\ &\ll T^{11/4-51/1168+\varepsilon} \\ &\ll T^{11/4-1/336+\varepsilon}. \end{aligned} \tag{3.32}$$

By Hölder's inequality, we have

$$\begin{aligned}\int_T^{2T} |R_1|^6 |R_2| \mathrm{d}x &\ll \left(\int_T^{2T} (|R_1|^6)^{7/6} \mathrm{d}x\right)^{6/7} \left(\int_T^{2T} |R_2|^7 \mathrm{d}x\right)^{1/7} \\ &\ll \left(\int_T^{2T} |R_1|^7 \mathrm{d}x\right)^{6/7} \left(\int_T^{2T} |R_2|^7 \mathrm{d}x\right)^{1/7} \\ &\ll (T^{11/4})^{6/7} \cdot (T^{11/4-51/1168+\varepsilon})^{1/7} \\ &\ll T^{11/4-1/336+\varepsilon}. \end{aligned} \tag{3.33}$$

Now Proposition 3.2 follows from (3.32) and (3.33).

From Proposition 3.1 and Proposition 3.2, we get the seventh-power moment of $\Delta(x)$.

Acknowledgement

The author would like to express the most sincere gratitude to Professor Wenguang Zhai for his valuable advice and constant encouragement.

References

[1] K. Corrádi and I. Kátai, *A comment on K. S. Gangadharan's paper entitled "Two classical lattice point problems"*, Magyar Tud. Akad. Mat. Fiz. Oszt. Kozl. 17 (1967) 89-97.

[2] J. L. Hafner, *New omega theorems for two classical lattice point problems*, Invent. Math. 63 (1981) 181-186.

[3] G. H. Hardy, *On Dirichlet's divisor problem*, Proc. London Math. Soc. (2) 15 (1916) 1-25.

Exponential sums with a difference, Proc. London Math. Soc. (3) 61 (1990) 227-250.

[4] D. R. Heath-Brown, *The distribution and moments of the error term in the Dirichlet divisor problem*, Acta Arith. 60 (1992) 389-415.

66 (1993), 279-301.

[5] M. N. Huxley, *Exponential sums and lattice points III*, Proc. London Math. Soc. (3) 87 (2003) 591-609.

[6] A. Ivić, *Large values of the error term in the divisor problem*, Invent. Math. 71 (1983) 513-520.

[7] A. Ivić, *Lectures on mean values of the Riemann zeta-function*, Tata Institute of Fundamental Research Lectures on Mathematics and Physics, Vol. 82, Bombay, 1991.

[8] A. Ivić and P. Sargos, *On the higher power moments of the error term in the divisor problem*, Illinois J. Math. 51 (2007) 353-377.

[9] K. L. Kong, *Some mean value theorems for certain error terms in analytic number theory*, Master degree Thesis, The University of Hong Kong, 2014.

[10] E. Krätzel, *Lattice points*, Deutsch. Verlag Wiss., Berlin, 1988.

[11] I. I. Piatetski-Shapiro, *On a variant of Waring-Goldbachs problem*, Mat. Sb. 30 (72) (1) (1952), 105-120 (in Russian).

[12] O. Robert and P. Sargos, *Three-dimensional exponential sums with monomials*, J. Reine Angew. Math. 591 (2006) 1-20.

[13] B. I. Segal, *On a theorem analogous to Warings theorem*, Dokl. Akad. Nauk SSSR (N. S.) 2 (1933), 47-49 (in Russian).

[14] K. C. Tong, *On divisor problem III*, Ada Math. Sinica 6 (1956) 515-541.

[15] K. M. Tsang, *Higher-power moments of* $\Delta(x), E(t)$ *and* $P(x)$, Proc. London Math. Soc. (3) 65 (1992) 65-84.

[16] G. F. Voronoï, *Sur une fonction transcendante et ses applications à la sommation de quelques séries*, Ann. École Normale (3) 21 (1904) 207-268, 459-534.

[17] J. Wang, *On the sixth-power moment of* $\Delta(x)$ *(in Chinese)*, Master degree Thesis, Shandong Normal University, 2010.

[18] W. G. Zhai, *On higher-power moments of* $\Delta(x)$, Acta Arith. 112 (2004) 367-395.

[19] W. G. Zhai, *On higher-power moments of* $\Delta(x)$ *(II)*, Acta Arith. 114 (2004) 35-54.

[20] W. G. Zhai, *On higher-power moments of* $\Delta(x)$ *(III)*, Acta Arith. 118 (2005) 263-281.

[21] D. Y. Zhang and W. G. Zhai, *On the fifth-power moment of* $\Delta(x)$, International Journal of Number Theory, Vol. 7, No. 1 (2011) 71-86.